

Math 474 - Homework # 4  
Random Variables, Expected Value, Games

1. The following game is called Chuck-a-luck. It works as follows. You pick a number out of 1, 2, 3, 4, 5, or 6 and bet \$1 on that number. Three giant 6-sided dice are then rolled in a spinning cage. You then win \$1 for every time that your number appears on the dice. But you lose your \$1 if your number doesn't appear at all. For example, suppose that you pick the number 1 as your number. Suppose that the dice show 1, 5, 1. Then you win \$2. If the dice showed 3, 1, 6 then you would win \$1. If the dice showed 3, 2, 6, then you would lose your \$1 bet.
  - (a) Let  $X$  denote the amount of money lost or won. Let  $p(i) = P(X = i)$  be the probability function for  $X$ . Calculate  $p(-1)$ ,  $p(1)$ ,  $p(2)$ ,  $p(3)$ .
  - (b) Draw a picture of  $p$ .
  - (c) Draw a picture of the cumulative distribution function  $F$  where  $F(i) = P(X \leq i)$ .
  - (d) What is the expected value of this game?
  
2. Consider the following experiment. Suppose we roll an 4-sided dice continually. We don't stop until a 3 is rolled.
  - (a) What is a sample space  $S$  and a probability function  $P$  for such an experiment? Verify that you have a probability space.
  - (b) Let  $A$  be the event that a 3 is rolled on the 3rd roll. Calculate  $P(A)$ .
  - (c) Let  $B$  be the event that a 3 is rolled within the first 3 rolls. Calculate  $P(B)$ .
  - (d) Suppose someone says this before the experiment starts: If a 3 is rolled within the first 3 rolls then I will pay you \$5, but if it doesn't then you have to pay me \$6. Do you take the bet?

3. Suppose there is a 4-sided die, but it isn't fair. Through experimentation you discover that a 1 is rolled approximately twice every eight rolls, a 2 is rolled approximately once every eight rolls, a 3 is rolled approximately three times every eight rolls, and a 4 is rolled approximately twice every eight rolls.
  - (a) Suppose you win \$2 for every 1 or 2 that is rolled, but lose \$1 for every 3 or 4 that is rolled. What is the expected value of such a game?
  - (b) What should you win or lose so that the game becomes "fair?" That is, so that the expected value is 0.
  
4. Suppose from a standard 52-card deck you are given the following five cards:  $4\heartsuit$ ,  $10\heartsuit$ ,  $Q\heartsuit$ ,  $3\spadesuit$ ,  $2\clubsuit$ . Suppose you now discard the  $3\spadesuit$  and  $2\clubsuit$  from your hand (but keep the other three cards) and ask for two more cards.
  - (a) What are the odds you get two more hearts so you have a flush?
  - (b) What are the odds you get one heart and a different suit?
  - (c) What are the odds you get exactly one more queen?
  - (d) What are the odds you get exactly two more queens?
  - (e) What are the odds that you get at least one more queen?
  - (f) Suppose someone says: If you get a flush I'll pay you \$500. But if you don't you have to pay me \$20. Do you take the bet?
  
5. Suppose that you flip a coin continually until a head occurs. Suppose that someone says: If you don't get a heads until you roll at least 3 tails then I'll pay you \$5. But if a heads occurs in the first three rolls then you must pay me \$1. Do you take the bet?